


Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

1. (Previously Presented) A compound of the formula (I)




wherein:

R_3 - R_5 are each independently selected from among hydrogen, C_{1-6} alkyls, C_{3-12} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} alkenyls, C_{3-12} branched alkenyls, C_{1-6} alkynyls, C_{3-12} branched alkynyls, C_{1-6} heteroalkyls, substituted C_{1-6} hetero-alkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys;

R_6 is OH, NH-aryl, NH-aralkyl, or NH- C_{1-12} alkyl,

w is 1 or 2;

Q_a is H or a residue of the formula:



wherein:


R_1 is a polyalkylene oxide;

Y_1 is O, S or NR_5 ; and

q is 0 or a positive integer;

d is 0 or 1; and

Q_b is H or a residue of the formula:



wherein:









R_2 is a polyalkylene oxide;






Y_2 is O, S or NR_5 ; and

s is 0 or a positive integer;

e is 0 or 1;

wherein L_{1-2} are independently selected from the group consisting of amino acids and





wherein:

R_{25} - R_{30} are independently selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{2-6} alkenyls, C_{2-6} alkynyls, C_{3-19} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{2-6} substituted alkenyls, C_{2-6} substituted alkynyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} heteroalkyls, substituted C_{1-6} hetero-alkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys;

R_{31} is selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{2-6} alkenyls, C_{2-6} alkynyls, C_{3-19} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{2-6} substituted alkenyls, C_{2-6} substituted alkynyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} heteroalkyls, substituted C_{1-6} heteroalkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys, NO_2 , haloalkyl and halogen;


t and y are individually selected positive integers, and

v is 0 or 1;

provided that Q_a and Q_b are both not simultaneously H.




3. (Currently Amended) The compound of claim 1 wherein R_2 further comprises a capping group J selected from the group consisting of OH, NH_2 , SH, CO_2H , C_{1-6} alkyl moieties, and a compound of the formula:




4. (Currently Amended) A compound of claim 2 of the formula:

(i)-R₁-(i)



wherein

(i) is:



wherein:

Y_1 is O;

L_1 is a hydrolysis-resistant bifunctional linker;

R_3 and R_4 are each independently hydrogen or CH_3 ;


R_6 is OH or NH-aryl;

q is 0-2; and

w is 1.


5. (Currently Amended) A compound of claim 3 ~~2~~ of the formula:

(ii)- R_2 -(ii)



wherein

(ii) is:



wherein:

Y_2 is O;

L_2 is a hydrolysis resistant bifunctional linker

R_3 and R_4 are each independently hydrogen or CH_3 ;

R_6 is OH or NH-aryl;

s is 0-2; and

w is 1.

6. (Original) The compound of claim 1 wherein:

Y_1 and Y_2 are independently O;

R_3 and R_4 are each independently hydrogen or CH_3 ;


R_6 is OH or NH-aryl;

q and s are independently 0-2; and

w is 1.

7. (Cancelled)

8. (Currently Amended) The compound of claim 1 ~~7~~ wherein the amino acid ~~residue~~ is selected from the group consisting of alanine, valine, leucine, isoleucine, glycine, serine, threonine, methionine, cysteine, phenylalanine, tyrosine, tryptophan, aspartic acid, glutamic acid, lysine, arginine, histidine and proline.
9. (Currently Amended) The compound of claim 1, wherein R_1 and R_2 independently comprise a linear, ~~terminally branched~~ or multi-armed polyalkylene oxide.
10. (Previously Presented) The compound of claim 9, wherein said polyalkylene oxide comprises polyethylene glycol.
11. (Currently Amended) The compound of claim 9, wherein said linear polyalkylene oxide ~~residue~~ is selected from the group consisting of:



wherein

A is a capping group;

R_7 is selected from that which defines R_3 , and

x is the degree of polymerization.

12. (Currently Amended) The compound of claim 9 ~~11~~ wherein said polyalkylene oxide ~~residue~~ has a total number average molecular weight of from about 5,000 to about 100,000 daltons.
13. (Currently Amended) The compound of claim 9 ~~11~~ wherein said polyalkylene oxide ~~residue~~ has a total number average molecular weight of from about 10,000 to about 80,000 daltons.

14. (Currently Amended) The compound of claim 9 ~~14~~ wherein said polyalkylene oxide residue has a total number average molecular weight of from about 20,000 to about 40,000 daltons.





15. (Cancelled)

16. (Currently Amended) The compound of claim 15, wherein said polyalkylene oxide residue comprises polyethylene glycol.


17. (Original) The compound of claim 16, wherein said polyethylene glycol has a number average molecular weight of from about 2,000 to about 100,000 daltons.

18. (Original) The compound of claim 16, wherein said polyethylene glycol has a number average molecular weight of from about 20,000 to about 40,000 daltons.

19. (Currently Amended) The compound of claim 1 9, selected from the group consisting of:



and



wherein

(a) is an integer of from about 1 to about 5;

X is O, NR₈, S, SO or SO₂; where R₈ is H, C₁₋₈ alkyl, C₁₋₈ branched alkyl, C₁₋₈ substituted alkyl, aryl or aralkyl;


(m) is 0 or 1;

(p) is a positive integer;

D is a moiety of the formula V_a or V_b,


wherein

V_a is a moiety of the formula:




; and

V_b is a moiety of the formula:



; and


mPEG is





wherein x is an integer from about 10 to about 2,300, and has a number average molecular weight of from about 2,000 to about 100,000 daltons.


20. (Original) The compound of claim 19, wherein mPEG has a number average molecular weight of from about 20,000 to about 40,000 daltons.

21. (Previously Amended) The compound of claim 1, selected from the group consisting of the formulas:





and



wherein,

m is 0-4;

z is 0 or 1;

L₄ is the same as that which defines L₁₋₂;

D is a moiety of the formula V_a or V_b;

R₁' is is $[[=]]$

$-(\text{CH}_2\text{CH}_2\text{O})_x-[[;]]_z$

$-(\text{CH}_2\text{CH}_2\text{O})_x-\text{CH}_2\text{C}(\text{O})-[[;]]_z$


$-(\text{CH}_2\text{CH}_2\text{O})_x-\text{CH}_2\text{CH}_2\text{NR}_7\text{-or and}$

$-(\text{CH}_2\text{CH}_2\text{O})_x-\text{CH}_2\text{CH}_2\text{SH}-[[;]]_z$

wherein


x is a positive integer;

V_a is a moiety of the formula:



; and


V_b is a moiety of the formula:











22. (Currently Amended) The compound of claim 21, wherein x is a positive integer such that the poly polymeric portion has a number average molecular weight of from about 2,000 to about 100,000 daltons.

23. (Currently Amended) The compound of claim 21, wherein x is a positive integer such that the poly polymeric portion has a number average molecular weight of from about 20,000 to about 40,000 daltons.

24. (Currently Amended) A compound selected from the group consisting of:








wherein:

PEG is



mPEG is



(a) is an integer of from about 1 to about 5;


Z is O, NR₈, S, SO or SO₂; where R₈ is H, C₁₋₈ alkyl, C₁₋₈ branched alkyl, C₁₋₈ substituted alkyl, aryl or aralkyl;

(m) is 0 or 1;

(p) is a positive integer;

x is 10 to 2,300; and






V_a is a moiety of the formula:




wherein:

Y₁ is O;

L₁ is selected from the group consisting of amino acids and





wherein:

R_{25} - R_{30} are independently selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{2-6} alkenyls, C_{2-6} alkynyls, C_{3-19} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{2-6} substituted alkenyls, C_{2-6} substituted alkynyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} heteroalkyls, substituted C_{1-6} hetero-alkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys;

R_{31} is selected from the group consisting of hydrogen, C_{1-6} alkyls,

C₂₋₆ alkenyls, C₂₋₆ alkynyls, C₃₋₁₉ branched alkyls, C₃₋₈ cycloalkyls, C₁₋₆ substituted alkyls, C₂₋₆ substituted alkenyls, C₂₋₆ substituted alkynyls, C₃₋₈ substituted cycloalkyls, aryls, substituted aryls, aralkyls, C₁₋₆ heteroalkyls, substituted C₁₋₆ heteroalkyls, C₁₋₆ alkoxyalkyl, phenoxyalkyl and C₁₋₆ heteroalkoxys, NO₂, haloalkyl and halogen;

t and y are individually selected positive integers, and

v is 0 or 1;

R₃ and R₄ are each independently hydrogen or CH₃;




R₆ is OH or NH-aryl;





q is 0-2;






d is 0 or 1; and





w is 1.


25. (Currently Amended) A compound selected from the group consisting of:












wherein:

PEG is



mPEG is



(a) is an integer of from about 1 to about 5;


Z is O, NR₈, S, SO or SO₂; where R₈ is H, C₁₋₈ alkyl, C₁₋₈ branched alkyl, C₁₋₈ substituted alkyl, aryl or aralkyl;

(m) is 0 or 1;

(p) is a positive integer, from about 1 to about 6;

x is 10 to 2,300, and


V_b is:







wherein:

Y_2 is O;

L_2 is a bifunctional linker selected from the group consisting of amino acids and





wherein:

R_{25} - R_{30} are independently selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{2-6} alkenyls, C_{2-6} alkynyls, C_{3-19} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{2-6} substituted alkenyls, C_{2-6} substituted alkynyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} heteroalkyls, substituted C_{1-6} hetero-alkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys;

R_{31} is selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{2-6} alkenyls, C_{2-6} alkynyls, C_{3-19} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{2-6} substituted alkenyls, C_{2-6} substituted alkynyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} heteroalkyls, substituted C_{1-6} heteroalkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys, NO_2 , haloalkyl and halogen;

t and y are individually selected positive integers, and

v is 0 or 1;

R_3 and R_4 are each independently hydrogen or CH_3 ;




R_6 is OH or NH-aryl;



s is 0-2;

e is 0 or 1; and


w is 1.

26. (Previously Presented) A compound of claim 1 having the formula:






and




wherein

 V_a is a moiety of the formula:




; and

V_b is a moiety of the formula:



27. (Withdrawn) A process for preparing a conjugate of claim 1 comprising, reacting a vancomycin compound of the formula:



wherein

R_3 and R_4 are independently selected from the group consisting of hydrogen, C_{1-6} alkyls, C_{3-12} branched alkyls, C_{3-8} cycloalkyls, C_{1-6} substituted alkyls, C_{3-8} substituted cycloalkyls, aryls, substituted aryls, aralkyls, C_{1-6} hetero-alkyls, substituted C_{1-6} hetero-alkyls, C_{1-6} alkoxyalkyl, phenoxyalkyl and C_{1-6} heteroalkoxys;

R_6 is OH, NH-aryl, NH-aralkyl, or NH- C_{1-12} alkyl; and

w is 1 or 2;

with a polymer residue containing at least one leaving group capable of reacting with the sugar amino group of said vancomycin compound in the presence of at least about a twenty-fold molar excess of triethylamine and a sufficient amount of dimethylformamide.

28. (Withdrawn) The process of claim 25 further comprising reacting said sugar amino conjugate with a second activated polymer residue containing at least one leaving group capable of reacting with the N-methyl-amino group of said conjugate in the presence of at least about a 5 fold molar excess of dimethylaminopyridine and a sufficient amount of a solvent mixture of dichloromethane and dimethylformamide.

29. (Withdrawn) The process of claim 26, wherein said solvent mixture comprises about equal parts dichloromethane and dimethylformamide.

30. (Withdrawn) A method of treating a vancomycin susceptible disease in a mammal comprising administering an effective amount of a compound of claim 1, to a mammal in need of such treatment, whereby, the compound of claim 1 undergoes degradation and releases vancomycin or a vancomycin derivative *in vivo*.
31. (Withdrawn) A method of treating a vancomycin susceptible disease in a mammal comprising administering an effective amount of a compound of claim 24, to a mammal in need of such treatment, whereby, the compound of claim 24 undergoes degradation and releases vancomycin or a vancomycin derivative *in vivo*.
32. (Withdrawn) A method of treating a vancomycin susceptible disease in a mammal comprising administering to a mammal in need of such treatment, an effective amount of a combination of vancomycin or a pharmaceutically acceptable salt, solvate or hydrate thereof, and a compound of claim 1.
33. (Original) A kit comprising in separate containers in a single package, pharmaceutical compositions for use in combination to treat a vancomycin susceptible disease which comprises in one container a therapeutically effective amount of vancomycin or a pharmaceutically acceptable salt, solvate or hydrate thereof in a pharmaceutically acceptable carrier and in a second container a therapeutically effective amount of a compound of claim 1 or a pharmaceutically acceptable salt, solvate or hydrate thereof in a pharmaceutically acceptable carrier.